

## Section 2.5 – Zeros of Polynomial Functions

The polynomial function  $f(x) = 63x^3 - 129x^2 + 28x + 20$  has  $x = \frac{5}{3}, -\frac{2}{7}, \frac{2}{3}$  as its zeros.

Notice that the NUMERATORS of these zeros (5, -2, and 2) are factors of the *constant term*, 20.

Also notice that the DENOMINATORS (3 and 7) are factors of the *leading coefficient*, 63.

### The Rational Zero Theorem

If  $f(x) = a_n x^n + \dots + a_1 x^1 + a_0$  has integer coefficients, then *every* rational zero of  $f(x)$  has the following form:

$$\frac{p}{q} = \frac{\text{factors of constant term}}{\text{factors of leading coefficient}}$$

Ex 1) Find the rational zeros of  $f(x) = x^3 + 2x^2 - 11x - 12$

1<sup>st</sup>: List the possible rational zeros

constant term's factors:

leading coefficient's factors:

So, the possible rational zeros are:

$$x = \underline{\hspace{2cm}} \rightarrow x = \underline{\hspace{2cm}}$$

2<sup>nd</sup>: Test these zeros using synthetic division

Try  $x = 1$

Try  $x = -1$

Since  $-1$  is a zero of  $f(x)$ , you can write the following:  $f(x) = \underline{\hspace{2cm}}$

Factor the trinomial and write the function in fully factored form:

$$f(x) = \underline{\hspace{2cm}} \text{ so the zeros are: } x = \underline{\hspace{1cm}}, \underline{\hspace{1cm}}, \underline{\hspace{1cm}}$$

## Section 2.5 – Zeros of Polynomial Functions

Ex 2) Find the rational zeros of  $f(x) = 2x^3 - 3x^2 - 8x - 3$

1<sup>st</sup>: List the possible rational zeros

The possible rational zeros are:

$x = \underline{\hspace{2cm}}$   $\rightarrow$   $x = \underline{\hspace{2cm}}$

2<sup>nd</sup>: Test these zeros using synthetic division

$f(x) = \underline{\hspace{2cm}}$

Factor the trinomial and write the function in fully factored form:

$f(x) = \underline{\hspace{2cm}}$  so the zeros are:  $x = \underline{\hspace{1cm}}, \underline{\hspace{1cm}}, \underline{\hspace{1cm}}$

## Section 2.5 – Zeros of Polynomial Functions

Ex 3) Use the following function  $f(x) = 2x^3 + 2x^2 - 8x - 8$  to:

Find possible zeros, pick some and test them using synthetic division, factor completely. Then state all zeros.

Ex 4) Use the following function  $f(x) = x^4 - x^3 + x^2 - 3x - 6$  to:

Find possible zeros, pick some and test them using synthetic division, factor completely. Then state all zeros.

## Section 2.5 – Zeros of Polynomial Functions

What about when not all the zeros are REAL?

**Conjugate Pairs Theorem:** Imaginary zeros always come in conjugate pairs.

Corollary to Conjugate Pairs Theorem:

A polynomial of odd degree must have at least one real zero.

Example: Find **all** zeros of the polynomial  $f(x) = 3x^4 + 5x^3 + 25x^2 + 45x - 18$

Example: Find **all** zeros of the polynomial  $f(x) = x^3 + 13x^2 + 57x + 85$

Homework: p. 179 #1, 3, 9, 11, 13, 23, 29, 65, 71

## Section 2.5 – Zeros of Polynomial Functions

There are a few tricks to help you get a zero with fewer guesses:

### Upper and Lower Bounds:

1. If you do synthetic division and get all positive numbers below the line, then the number you used is too high – go lower with your next guess!!
2. If you do synthetic division and get a pattern of positive, negative, positive, negative (or neg-pos-neg-pos) for your numbers below the line, then your guess is too low – go higher with your next guess!!

### Descartes' Rule of Signs:

1. If you look at  $f(x)$  and count the number of sign changes, that number (possibly less multiples of two) will give you the number of positive real zeros for  $f(x)$ .
2. If you look at  $f(-x)$  and count the number of sign changes, that number (possibly less multiples of two) will give you the number of negative real zeros for  $f(x)$ .